LISTA DE PUBLICAŢII REPREZENTATIVE

1. **Kiss E**, Popovic Z, Bedke J, Wang S, Bonrouhi M, Gretz N, Stettner P, Teupser D, Thiery J, Porubsky S, Adams J, Gröne HJ. Suppression of chronic damage in renal allografts by Liver X receptor (LXR) activation relevant contribution of macrophage LXRα. Am J Pathol. 2011;179(1):92-103. doi:10.1016/j.ajpath.2011.03.019. FI: 4,89
2. **Kiss E**, Kränzlin B, Wagenblaβ K, Bonrouhi M, Thiery J, Gröne E, Nordström V, Teupser D, Gretz N, Malle E, Gröne HJ. Lipid droplet accumulation is associated with an increase in hyperglycemia-induced renal damage: prevention by liver X receptors. Am J Pathol. 2013;182(3):727-41. doi: 10.1016/j.ajpath.2012.11.033. FI: 4,6
3. **Kiss E**, Gorgas K, Schlicksupp A, Groß D, Kins S, Kirsch J, Kuhse J. Biphasic Alteration of the Inhibitory Synapse Scaffold Protein Gephyrin in Early and Late Stages of an Alzheimer Disease Model. Am J Pathol. 2016;186(9):2279-91. doi: 10.1016/j.ajpath.2016.05.013. FI: 4,05
4. Hollnagel JO, Elzoheiry S, Gorgas K, Kins S, Beretta CA, Kirsch J, Kuhse J, Kann O, **Kiss E**. Early alterations in hippocampal perisomatic GABAergic synapses and network oscillations in a mouse model of Alzheimer's disease amyloidosis. PLoS One. 2019;14(1):e0209228. doi: 10.1371/journal.pone.0209228. FI: 2,76
5. **Kiss E**, Groeneweg F, Gorgas K, Schlicksupp A, Kins S, Kirsch J, Kuhse J. Amyloid-β Fosters p35/CDK5 Signaling Contributing to Changes of Inhibitory Synapses in Early Stages of Cerebral Amyloidosis. J Alzheimers Dis. 2020;74(4):1167-1187. doi: 10.3233/JAD-190976. IF:4,47
6. **Kiss E**, Kins S, Gorgas K, Orlik M, Fischer C, Endres K, Schlicksupp A, Kirsch J, Kuhse J. Artemisinin-treatment in pre-symptomatic APP-PS1 mice increases gephyrin phosphorylation at Ser270: a modification regulating postsynaptic GABAAR density. Biol Chem. 2021;403(1):73-87. doi: 10.1515/hsz-2021-0153. IF:3,46
7. **Kiss E**, Kins S, Zöller Y, Schilling S, Gorgas K, Groß D, Schlicksupp A, Rosner R, Kirsch J, Kuhse J. Artesunate restores the levels of inhibitory synapse proteins and reduces amyloid-β and C-terminal fragments (CTFs) of the amyloid precursor protein in an AD-mouse model. Mol Cell Neurosci. 2021 ;113:103624. doi: 10.1016/j.mcn.2021.103624. IF:3,18
8. Kuhse J, Groeneweg F, Kins S, Gorgas K, Nawrotzki R, Kirsch J, **Kiss E**. Loss of Extrasynaptic Inhibitory Glycine Receptors in the Hippocampus of an AD Mouse Model Is Restored by Treatment with Artesunate. Int J Mol Sci. 2023;24(5):4623. doi: 10.3390/ijms24054623. IF:5,6
9. **Kiss E**, Kins S, Gorgas K, Venczel Szakács KH, Kirsch J, Kuhse J. Another Use for a Proven Drug: Experimental Evidence for the Potential of Artemisinin and Its Derivatives to Treat Alzheimer's Disease. Int J Mol Sci. 2024 Apr 9;25(8):4165. doi: 10.3390/ijms25084165.
10. **Kiss E**, Kirsch J, Kuhse J, Kins S. Is the Activation of the Postsynaptic Ligand Gated Glycine- or $GABA_A$ Receptors Essential for the Receptor Clustering at Inhibitory Synapses? Biomedicines. 2025 Aug 5;13(8):1905. doi: 10.3390/biomedicines13081905. IF:4,9